

# MEM6810 Engineering Systems Modeling and Simulation



## 工程系统建模与仿真

Theory Analysis

### Lecture 7: Output Analysis I: Single Model

SHEN Haihui 沈海辉

Sino-US Global Logistics Institute  
Shanghai Jiao Tong University

 [shenhaihui.github.io/teaching/mem6810f](https://shenhaihui.github.io/teaching/mem6810f)  
 [shenhaihui@sjtu.edu.cn](mailto:shenhaihui@sjtu.edu.cn)

Spring 2023 (full-time)



上海交通大学  
SHANGHAI JIAO TONG UNIVERSITY

董浩云智能制造与服务管理研究院  
CY TUNG Institute of Intelligent Manufacturing and Service Management  
(中美物流研究院)  
(Sino-US Global Logistics Institute)



- 1 Introduction
  - ▶ Types of Simulations
- 2 Point and Interval Estimations
  - ▶ Basics
  - ▶ Specified Precision
  - ▶ Example
- 3 Terminating Simulation
  - ▶ Discrete Outputs
  - ▶ Continuous Outputs
- 4 Steady-State Simulation
  - ▶ Initialization Bias
  - ▶ Intelligent Initialization
  - ▶ Warm-up Period Deletion
  - ▶ Estimation with Multiple Replications
  - ▶ Estimation with Single Replication



- 1 Introduction
  - ▶ Types of Simulations
- 2 Point and Interval Estimations
  - ▶ Basics
  - ▶ Specified Precision
  - ▶ Example
- 3 Terminating Simulation
  - ▶ Discrete Outputs
  - ▶ Continuous Outputs
- 4 Steady-State Simulation
  - ▶ Initialization Bias
  - ▶ Intelligent Initialization
  - ▶ Warm-up Period Deletion
  - ▶ Estimation with Multiple Replications
  - ▶ Estimation with Single Replication



# Introduction

- Output analysis is the examination of data generated by a simulation.
- Output analysis is needed because the output data from a stochastic simulation exhibits random variability.

- Output analysis is the examination of data generated by a simulation.
- Output analysis is needed because the output data from a stochastic simulation exhibits random variability.
- Suppose the true performance of the *simulated system* is  $\theta$ .
  - The result from a set of simulation runs will be an estimator  $\hat{\theta}$ .
  - The precision of the estimator  $\hat{\theta}$  can be measured by the **standard error** of  $\hat{\theta}$  (i.e., estimator's standard deviation) or the width of a confidence interval for  $\theta$ .

- Output analysis is the examination of data generated by a simulation.
- Output analysis is needed because the output data from a stochastic simulation exhibits random variability.
- Suppose the true performance of the *simulated system* is  $\theta$ .
  - The result from a set of simulation runs will be an estimator  $\hat{\theta}$ .
  - The precision of the estimator  $\hat{\theta}$  can be measured by the **standard error** of  $\hat{\theta}$  (i.e., estimator's standard deviation) or the width of a confidence interval for  $\theta$ .
- The purpose of the statistical analysis:
  - Estimate the true performance  $\theta$ .
  - Control the estimation precision.

- Output analysis is the examination of data generated by a simulation.
- Output analysis is needed because the output data from a stochastic simulation exhibits random variability.
- Suppose the true performance of the *simulated system* is  $\theta$ .
  - The result from a set of simulation runs will be an estimator  $\hat{\theta}$ .
  - The precision of the estimator  $\hat{\theta}$  can be measured by the **standard error** of  $\hat{\theta}$  (i.e., estimator's standard deviation) or the width of a confidence interval for  $\theta$ .
- The purpose of the statistical analysis:
  - Estimate the true performance  $\theta$ .
  - Control the estimation precision.
- Types of simulation with regard to output analysis:
  - terminating vs. nonterminating.

- A **terminating simulation** is one that runs for some well-defined time duration  $T_E$ .
  - $E$  is a specified event (or set of events) that stops each simulation run (replication).
  - Simulation starts at time 0 under well-specified initial conditions, and ends at the stopping time  $T_E$ .
  - $T_E$  can be either a deterministic or random variable.



- A **terminating simulation** is one that runs for some well-defined time duration  $T_E$ .
  - $E$  is a specified event (or set of events) that stops each simulation run (replication).
  - Simulation starts at time 0 under well-specified initial conditions, and ends at the stopping time  $T_E$ .
  - $T_E$  can be either a deterministic or random variable.
- Example: A bank opens at 9 AM (*time 0*) with no customers present and 8 of the 11 tellers working (*initial conditions*), and closes at 5 PM (*time  $T_E = 8$  hours*).
  - $E = \{8 \text{ hours of simulated time have elapsed}\}$ .

- A **terminating simulation** is one that runs for some well-defined time duration  $T_E$ .
  - $E$  is a specified event (or set of events) that stops each simulation run (replication).
  - Simulation starts at time 0 under well-specified initial conditions, and ends at the stopping time  $T_E$ .
  - $T_E$  can be either a deterministic or random variable.
- Example: A bank opens at 9 AM (*time 0*) with no customers present and 8 of the 11 tellers working (*initial conditions*), and closes at 5 PM (*time  $T_E = 8$  hours*).
  - $E = \{8 \text{ hours of simulated time have elapsed}\}$ .
- It actually stops service when the last customer who entered before 5 PM has been served.
  - $E = \{\text{at least 8 hours of simulated time have elapsed and the system is empty}\} \implies T_E$  is a random variable.

- A **nonterminating simulation** is one that runs continuously and without a natural event  $E$  to stop the simulation run.
  - Initial conditions are defined by the analyst, but its effect *fades away* as simulation time increases.
  - Stopping time is conceptually infinite, and in practice it is determined by the analyst with certain statistical precision.

- A **nonterminating simulation** is one that runs continuously and without a natural event  $E$  to stop the simulation run.
  - Initial conditions are defined by the analyst, but its effect *fades away* as simulation time increases.
  - Stopping time is conceptually infinite, and in practice it is determined by the analyst with certain statistical precision.
- Examples: Production line that runs 24/7, hospital emergency rooms, continuously operating computer networks, etc.

- A **nonterminating simulation** is one that runs continuously and without a natural event  $E$  to stop the simulation run.
  - Initial conditions are defined by the analyst, but its effect *fades away* as simulation time increases.
  - Stopping time is conceptually infinite, and in practice it is determined by the analyst with certain statistical precision.
- Examples: Production line that runs 24/7, hospital emergency rooms, continuously operating computer networks, etc.
- For a simulation model that is run in a nonterminating way and *has a steady-state (stationary) distribution*:
  - The objective is often to study the long-run, or steady-state, behavior of a system, which is not influenced by the initial conditions.
  - Such nonterminating simulation is also called *steady-state simulation*.

- 1 Introduction
  - ▶ Types of Simulations
- 2 Point and Interval Estimations
  - ▶ Basics
  - ▶ Specified Precision
  - ▶ Example
- 3 Terminating Simulation
  - ▶ Discrete Outputs
  - ▶ Continuous Outputs
- 4 Steady-State Simulation
  - ▶ Initialization Bias
  - ▶ Intelligent Initialization
  - ▶ Warm-up Period Deletion
  - ▶ Estimation with Multiple Replications
  - ▶ Estimation with Single Replication



- Suppose we want to estimate  $\theta = \mathbb{E}[X]$  based on the iid sample  $\{X_1, \dots, X_n\}$ .

- Suppose we want to estimate  $\theta = \mathbb{E}[X]$  based on the iid sample  $\{X_1, \dots, X_n\}$ .
- The point estimator (a single random variable) is

$$\hat{\theta} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i.$$



- Suppose we want to estimate  $\theta = \mathbb{E}[X]$  based on the iid sample  $\{X_1, \dots, X_n\}$ .
- The point estimator (a single random variable) is

$$\hat{\theta} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i.$$

- How good is this estimator?
  - Unbiased:  $\mathbb{E}[\hat{\theta}] = \theta$ .
  - Consistent:  $\hat{\theta} \xrightarrow{a.s.} \theta$ , as  $n \rightarrow \infty$ .<sup>†</sup>

---

<sup>†</sup> Assume  $\mathbb{E}[|X|] < \infty$ , or, a stronger condition,  $\text{Var}(X) < \infty$ .



- Suppose we want to estimate  $\theta = \mathbb{E}[X]$  based on the iid sample  $\{X_1, \dots, X_n\}$ .
- The point estimator (a single random variable) is

$$\hat{\theta} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i.$$

- How good is this estimator?
  - Unbiased:  $\mathbb{E}[\hat{\theta}] = \theta$ .
  - Consistent:  $\hat{\theta} \xrightarrow{a.s.} \theta$ , as  $n \rightarrow \infty$ .<sup>†</sup>
- Point estimator says *nothing* about the estimation error for finite sample size  $n$ .
  - Small estimation error means high estimation precision.

---

<sup>†</sup> Assume  $\mathbb{E}[|X|] < \infty$ , or, a stronger condition,  $\text{Var}(X) < \infty$ .

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \sim \mathcal{N}(0, 1).$$

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \sim \mathcal{N}(0, 1).$$

- If  $X$  follows arbitrary distribution and  $\sigma^2 = \text{Var}(X) \in (0, \infty)$ , then by Central Limit Theorem,

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{as } n \rightarrow \infty. \quad (1)$$

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \sim \mathcal{N}(0, 1).$$

- If  $X$  follows arbitrary distribution and  $\sigma^2 = \text{Var}(X) \in (0, \infty)$ , then by Central Limit Theorem,

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{as } n \rightarrow \infty. \quad (1)$$

- $\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \sim \mathcal{N}(0, 1)$  approximately when  $n$  is large.

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \sim \mathcal{N}(0, 1).$$

- If  $X$  follows arbitrary distribution and  $\sigma^2 = \text{Var}(X) \in (0, \infty)$ , then by Central Limit Theorem,

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{as } n \rightarrow \infty. \quad (1)$$

- $\sqrt{n} \left( \frac{\hat{\theta} - \theta}{\sigma} \right) \sim \mathcal{N}(0, 1)$  approximately when  $n$  is large.
- $\sigma^2$  is typically unknown, and we substitute it by the sample variance

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2.$$

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \sim t_{n-1}, \quad (2)$$

where  $t_p$  denotes  $t$  distribution with  $p$  degrees of freedom.

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \sim t_{n-1}, \quad (2)$$

where  $t_p$  denotes  $t$  distribution with  $p$  degrees of freedom.

- If  $X$  follows arbitrary distribution and  $\sigma^2 = \text{Var}(X) \in (0, \infty)$ , then by Equation (1) and the fact that  $\frac{\sigma}{S} \xrightarrow{a.s.} 1$  as  $n \rightarrow \infty$ ,

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{as } n \rightarrow \infty. \quad (3)$$



- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \sim t_{n-1}, \quad (2)$$

where  $t_p$  denotes  $t$  distribution with  $p$  degrees of freedom.

- If  $X$  follows arbitrary distribution and  $\sigma^2 = \text{Var}(X) \in (0, \infty)$ , then by Equation (1) and the fact that  $\frac{\sigma}{S} \xrightarrow{a.s.} 1$  as  $n \rightarrow \infty$ ,

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{as } n \rightarrow \infty. \quad (3)$$

- $\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \sim \mathcal{N}(0, 1)$  approximately when  $n$  is large.

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , then

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \sim t_{n-1}, \quad (2)$$

where  $t_p$  denotes  $t$  distribution with  $p$  degrees of freedom.

- If  $X$  follows arbitrary distribution and  $\sigma^2 = \text{Var}(X) \in (0, \infty)$ , then by Equation (1) and the fact that  $\frac{\sigma}{S} \xrightarrow{a.s.} 1$  as  $n \rightarrow \infty$ ,

$$\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{as } n \rightarrow \infty. \quad (3)$$

- $\sqrt{n} \left( \frac{\hat{\theta} - \theta}{S} \right) \sim \mathcal{N}(0, 1)$  approximately when  $n$  is large.
- Results (2) and (3) are the basis of the confidence interval estimation for  $\theta$ .

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , where  $\theta$  and  $\sigma$  are unknown, then a  $1 - \alpha$  confidence interval (CI) for  $\theta$  is

$$\left[ \hat{\theta} - t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (4)$$

where  $t_{n-1, 1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $t_{n-1}$  distribution.

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , where  $\theta$  and  $\sigma$  are unknown, then a  $1 - \alpha$  confidence interval (CI) for  $\theta$  is

$$\left[ \hat{\theta} - t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (4)$$

where  $t_{n-1, 1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $t_{n-1}$  distribution.

Proof.

$$\mathbb{P} \left\{ \theta \in \left[ \hat{\theta} - t_{n-1, 1-\alpha/2} S / \sqrt{n}, \hat{\theta} + t_{n-1, 1-\alpha/2} S / \sqrt{n} \right] \right\}$$

- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , where  $\theta$  and  $\sigma$  are unknown, then a  $1 - \alpha$  confidence interval (CI) for  $\theta$  is

$$\left[ \hat{\theta} - t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (4)$$

where  $t_{n-1, 1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $t_{n-1}$  distribution.

Proof.

$$\begin{aligned} & \mathbb{P} \left\{ \theta \in \left[ \hat{\theta} - t_{n-1, 1-\alpha/2} S / \sqrt{n}, \hat{\theta} + t_{n-1, 1-\alpha/2} S / \sqrt{n} \right] \right\} \\ &= \mathbb{P} \left\{ |\theta - \hat{\theta}| \leq t_{n-1, 1-\alpha/2} S / \sqrt{n} \right\} \end{aligned}$$



- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , where  $\theta$  and  $\sigma$  are unknown, then a  $1 - \alpha$  confidence interval (CI) for  $\theta$  is

$$\left[ \hat{\theta} - t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (4)$$

where  $t_{n-1, 1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $t_{n-1}$  distribution.

Proof.

$$\begin{aligned} & \mathbb{P} \left\{ \theta \in \left[ \hat{\theta} - t_{n-1, 1-\alpha/2} S/\sqrt{n}, \hat{\theta} + t_{n-1, 1-\alpha/2} S/\sqrt{n} \right] \right\} \\ &= \mathbb{P} \left\{ |\theta - \hat{\theta}| \leq t_{n-1, 1-\alpha/2} S/\sqrt{n} \right\} \\ &= \mathbb{P} \left\{ \left| \frac{\theta - \hat{\theta}}{S/\sqrt{n}} \right| \leq t_{n-1, 1-\alpha/2} \right\} \end{aligned}$$



- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , where  $\theta$  and  $\sigma$  are unknown, then a  $1 - \alpha$  confidence interval (CI) for  $\theta$  is

$$\left[ \hat{\theta} - t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (4)$$

where  $t_{n-1, 1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $t_{n-1}$  distribution.

Proof.

$$\begin{aligned} & \mathbb{P} \left\{ \theta \in \left[ \hat{\theta} - t_{n-1, 1-\alpha/2} S / \sqrt{n}, \hat{\theta} + t_{n-1, 1-\alpha/2} S / \sqrt{n} \right] \right\} \\ &= \mathbb{P} \left\{ |\theta - \hat{\theta}| \leq t_{n-1, 1-\alpha/2} S / \sqrt{n} \right\} \\ &= \mathbb{P} \left\{ \left| \frac{\theta - \hat{\theta}}{S / \sqrt{n}} \right| \leq t_{n-1, 1-\alpha/2} \right\} = 1 - \alpha, \end{aligned}$$



- If  $X \sim \mathcal{N}(\theta, \sigma^2)$ , where  $\theta$  and  $\sigma$  are unknown, then a  $1 - \alpha$  confidence interval (CI) for  $\theta$  is

$$\left[ \hat{\theta} - t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (4)$$

where  $t_{n-1, 1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $t_{n-1}$  distribution.

Proof.

$$\begin{aligned} & \mathbb{P} \left\{ \theta \in \left[ \hat{\theta} - t_{n-1, 1-\alpha/2} S / \sqrt{n}, \hat{\theta} + t_{n-1, 1-\alpha/2} S / \sqrt{n} \right] \right\} \\ &= \mathbb{P} \left\{ |\theta - \hat{\theta}| \leq t_{n-1, 1-\alpha/2} S / \sqrt{n} \right\} \\ &= \mathbb{P} \left\{ \left| \frac{\theta - \hat{\theta}}{S / \sqrt{n}} \right| \leq t_{n-1, 1-\alpha/2} \right\} = 1 - \alpha, \end{aligned}$$


where the last equality is due to (2) and the symmetry of  $t$  distribution.






- The interpretation of CI:
  - If one constructs a very large number of independent  $1 - \alpha$  CIs, each based on  $n$  observations, the proportion of CIs that actually contain (cover)  $\theta$  should be  $1 - \alpha$ .

- The interpretation of CI:
  - If one constructs a very large number of independent  $1 - \alpha$  CIs, each based on  $n$  observations, the proportion of CIs that actually contain (cover)  $\theta$  should be  $1 - \alpha$ .




- The interpretation of CI:
  - If one constructs a very large number of independent  $1 - \alpha$  CIs, each based on  $n$  observations, the proportion of CIs that actually contain (cover)  $\theta$  should be  $1 - \alpha$ .



- **Caution:** There is nothing probabilistic about a single CI **after** the data have been observed and the interval's endpoints have been given numerical values, e.g.,  $[1.1, 2.4]$ .

- The interpretation of CI:
  - If one constructs a very large number of independent  $1 - \alpha$  CIs, each based on  $n$  observations, the proportion of CIs that actually contain (cover)  $\theta$  should be  $1 - \alpha$ .



- **Caution:** There is nothing probabilistic about a single CI **after** the data have been observed and the interval's endpoints have been given numerical values, e.g.,  $[1.1, 2.4]$ .
- Try it out! <http://www.rossmanchance.com/applets/ConfSim.html>



- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).

- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).
- The interpretation is the same as before.

- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).
- The interpretation is the same as before.
- In practice, people may also use (4) as approximate CI even when  $X$  does not follow a normal distribution.



- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).
- The interpretation is the same as before.
- In practice, people may also use (4) as approximate CI even when  $X$  does not follow a normal distribution.
  - Both (4) and (5) are approximation for finite  $n$  when  $X$  is non-normal.

- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).
- The interpretation is the same as before.
- In practice, people may also use (4) as approximate CI even when  $X$  does not follow a normal distribution.
  - Both (4) and (5) are approximation for finite  $n$  when  $X$  is non-normal.
  - $t_{n-1, 1-\alpha/2} > z_{1-\alpha/2}$ , so CI (4) will be wider than CI (5).

- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).
- The interpretation is the same as before.
- In practice, people may also use (4) as approximate CI even when  $X$  does not follow a normal distribution.
  - Both (4) and (5) are approximation for finite  $n$  when  $X$  is non-normal.
  - $t_{n-1, 1-\alpha/2} > z_{1-\alpha/2}$ , so CI (4) will be wider than CI (5).
  - CI (4) generally has coverage closer to the desired level  $1 - \alpha$ .

- If  $X$  follows arbitrary distribution,  $\theta$  and  $\sigma^2 = \text{Var}(X)$  are unknown, and  $0 < \sigma^2 < \infty$ , then an approximate  $1 - \alpha$  CI for  $\theta$  with large  $n$  is

$$\left[ \hat{\theta} - z_{1-\alpha/2} \frac{S}{\sqrt{n}}, \hat{\theta} + z_{1-\alpha/2} \frac{S}{\sqrt{n}} \right], \quad (5)$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ -quantile of  $\mathcal{N}(0, 1)$ .

- The proof is similar as before by using (3), instead of (2).
- The interpretation is the same as before.
- In practice, people may also use (4) as approximate CI even when  $X$  does not follow a normal distribution.
  - Both (4) and (5) are approximation for finite  $n$  when  $X$  is non-normal.
  - $t_{n-1, 1-\alpha/2} > z_{1-\alpha/2}$ , so CI (4) will be wider than CI (5).
  - CI (4) generally has coverage closer to the desired level  $1 - \alpha$ .
  - $t_{n-1, 1-\alpha/2} \rightarrow z_{1-\alpha/2}$  as  $n \rightarrow \infty$ .

- For CI (4), the half-length under  $1 - \alpha$  confidence level is

$$H = t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- For CI (4), the half-length under  $1 - \alpha$  confidence level is

$$H = t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- For CI (5), the half-length under  $1 - \alpha$  confidence level is

$$H = z_{1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- For CI (4), the half-length under  $1 - \alpha$  confidence level is

$$H = t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- For CI (5), the half-length under  $1 - \alpha$  confidence level is

$$H = z_{1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- Half-length  $H$  presents the precision (or error) of the estimation for  $\theta$ .

- For CI (4), the half-length under  $1 - \alpha$  confidence level is

$$H = t_{n-1, 1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- For CI (5), the half-length under  $1 - \alpha$  confidence level is

$$H = z_{1-\alpha/2} \frac{S}{\sqrt{n}}.$$

- Half-length  $H$  presents the precision (or error) of the estimation for  $\theta$ .
- We want  $H$  to be small enough for our decision making, say,  $H \leq \epsilon$ , under  $1 - \alpha$  confidence level.



- Usually we take an initial sample of size  $n_0$  to get an estimate of  $\sigma^2$ , say  $S_0^2$ .
  - Assume that the estimate of  $\sigma^2$  will not change (appreciably) from  $S_0^2$  as the sample size increases.

- Usually we take an initial sample of size  $n_0$  to get an estimate of  $\sigma^2$ , say  $S_0^2$ .
  - Assume that the estimate of  $\sigma^2$  will not change (appreciably) from  $S_0^2$  as the sample size increases.
- For CI (4), an approximate expression for the total sample size required to make  $H \leq \epsilon$  is given by

$$n^* = \min \left\{ n \geq n_0 : t_{n-1, 1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\}.$$

- Usually we take an initial sample of size  $n_0$  to get an estimate of  $\sigma^2$ , say  $S_0^2$ .
  - Assume that the estimate of  $\sigma^2$  will not change (appreciably) from  $S_0^2$  as the sample size increases.
- For CI (4), an approximate expression for the total sample size required to make  $H \leq \epsilon$  is given by

$$n^* = \min \left\{ n \geq n_0 : t_{n-1, 1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\}.$$

- For CI (5), an approximate expression is given by

$$n^* = \min \left\{ n \geq n_0 : z_{1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\} = \left\lceil \left( \frac{z_{1-\alpha/2} S_0}{\epsilon} \right)^2 \right\rceil. \quad (6)$$

- Usually we take an initial sample of size  $n_0$  to get an estimate of  $\sigma^2$ , say  $S_0^2$ .
  - Assume that the estimate of  $\sigma^2$  will not change (appreciably) from  $S_0^2$  as the sample size increases.

- For CI (4), an approximate expression for the total sample size required to make  $H \leq \epsilon$  is given by

$$n^* = \min \left\{ n \geq n_0 : t_{n-1, 1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\}.$$

- For CI (5), an approximate expression is given by

$$n^* = \min \left\{ n \geq n_0 : z_{1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\} = \left\lceil \left( \frac{z_{1-\alpha/2} S_0}{\epsilon} \right)^2 \right\rceil. \quad (6)$$

- For simplicity, people sometimes use (6), regardless of the distribution of  $X$ .

- Usually we take an initial sample of size  $n_0$  to get an estimate of  $\sigma^2$ , say  $S_0^2$ .
  - Assume that the estimate of  $\sigma^2$  will not change (appreciably) from  $S_0^2$  as the sample size increases.

- For CI (4), an approximate expression for the total sample size required to make  $H \leq \epsilon$  is given by

$$n^* = \min \left\{ n \geq n_0 : t_{n-1, 1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\}.$$

- For CI (5), an approximate expression is given by

$$n^* = \min \left\{ n \geq n_0 : z_{1-\alpha/2} \frac{S_0}{\sqrt{n}} \leq \epsilon \right\} = \left\lceil \left( \frac{z_{1-\alpha/2} S_0}{\epsilon} \right)^2 \right\rceil. \quad (6)$$

- For simplicity, people sometimes use (6), regardless of the distribution of  $X$ .
- Take  $n^* - n_0$  additional sample points, or start over and take a sample of size  $n^*$ , to form the  $1 - \alpha$  CI (with new  $S$ ).

- Suppose an iid sample is taken and the values are as follows:

79.919	3.081	0.062	1.961	5.845	0.941	0.878	3.371	2.157	7.579
3.027	6.505	0.021	0.013	0.123	0.624	5.380	3.148	7.078	23.960
6.769	59.899	1.192	34.760	5.009	0.590	1.928	0.300	0.002	0.543
18.387	0.141	43.565	24.420	0.433	7.004	31.764	1.005	1.147	0.219
144.695	2.663	17.967	0.091	9.003	3.217	14.382	1.008	2.336	4.562

- Suppose an iid sample is taken and the values are as follows:

79.919	3.081	0.062	1.961	5.845	0.941	0.878	3.371	2.157	7.579
3.027	6.505	0.021	0.013	0.123	0.624	5.380	3.148	7.078	23.960
6.769	59.899	1.192	34.760	5.009	0.590	1.928	0.300	0.002	0.543
18.387	0.141	43.565	24.420	0.433	7.004	31.764	1.005	1.147	0.219
144.695	2.663	17.967	0.091	9.003	3.217	14.382	1.008	2.336	4.562

- Construct a 95% CI and a 99% CI for  $\theta = \mathbb{E}[X]$ .

- Suppose an iid sample is taken and the values are as follows:

79.919	3.081	0.062	1.961	5.845	0.941	0.878	3.371	2.157	7.579
3.027	6.505	0.021	0.013	0.123	0.624	5.380	3.148	7.078	23.960
6.769	59.899	1.192	34.760	5.009	0.590	1.928	0.300	0.002	0.543
18.387	0.141	43.565	24.420	0.433	7.004	31.764	1.005	1.147	0.219
144.695	2.663	17.967	0.091	9.003	3.217	14.382	1.008	2.336	4.562

- Construct a 95% CI and a 99% CI for  $\theta = \mathbb{E}[X]$ .

$n = 50$ ,  $\hat{\theta} = \bar{X} = 11.894$ ,  $S = 24.953$ . We use CI (4) and get  $t_{49, 0.975} = 2.010$ ,  $t_{49, 0.995} = 2.680$ . Then,

$$95\% \text{ CI: } 11.894 \pm 2.010 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 7.093 = [4.801, 18.987];$$

$$99\% \text{ CI: } 11.894 \pm 2.680 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 9.457 = [2.437, 21.351].$$





- Suppose an iid sample is taken and the values are as follows:

79.919	3.081	0.062	1.961	5.845	0.941	0.878	3.371	2.157	7.579
3.027	6.505	0.021	0.013	0.123	0.624	5.380	3.148	7.078	23.960
6.769	59.899	1.192	34.760	5.009	0.590	1.928	0.300	0.002	0.543
18.387	0.141	43.565	24.420	0.433	7.004	31.764	1.005	1.147	0.219
144.695	2.663	17.967	0.091	9.003	3.217	14.382	1.008	2.336	4.562

- Construct a 95% CI and a 99% CI for  $\theta = \mathbb{E}[X]$ .

$n = 50$ ,  $\hat{\theta} = \bar{X} = 11.894$ ,  $S = 24.953$ . We use CI (4) and get  $t_{49, 0.975} = 2.010$ ,  $t_{49, 0.995} = 2.680$ . Then,

$$95\% \text{ CI: } 11.894 \pm 2.010 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 7.093 = [4.801, 18.987];$$

$$99\% \text{ CI: } 11.894 \pm 2.680 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 9.457 = [2.437, 21.351].$$

- Want to make half-length  $H \leq 2$  under 95% confidence level.

- Suppose an iid sample is taken and the values are as follows:

79.919	3.081	0.062	1.961	5.845	0.941	0.878	3.371	2.157	7.579
3.027	6.505	0.021	0.013	0.123	0.624	5.380	3.148	7.078	23.960
6.769	59.899	1.192	34.760	5.009	0.590	1.928	0.300	0.002	0.543
18.387	0.141	43.565	24.420	0.433	7.004	31.764	1.005	1.147	0.219
144.695	2.663	17.967	0.091	9.003	3.217	14.382	1.008	2.336	4.562

- Construct a 95% CI and a 99% CI for  $\theta = \mathbb{E}[X]$ .

$n = 50$ ,  $\hat{\theta} = \bar{X} = 11.894$ ,  $S = 24.953$ . We use CI (4) and get  $t_{49, 0.975} = 2.010$ ,  $t_{49, 0.995} = 2.680$ . Then,

$$95\% \text{ CI: } 11.894 \pm 2.010 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 7.093 = [4.801, 18.987];$$

$$99\% \text{ CI: } 11.894 \pm 2.680 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 9.457 = [2.437, 21.351].$$

- Want to make half-length  $H \leq 2$  under 95% confidence level.

We use (6) and get  $z_{0.975} = 1.960$ ,  $S_0 = 24.953$ ,  $\epsilon = 2$ . Then,

$$n^* = \left\lceil \left( \frac{1.960 \times 24.953}{2} \right)^2 \right\rceil = \lceil 597.995 \rceil = 598.$$

- Suppose an iid sample is taken and the values are as follows:

79.919	3.081	0.062	1.961	5.845	0.941	0.878	3.371	2.157	7.579
3.027	6.505	0.021	0.013	0.123	0.624	5.380	3.148	7.078	23.960
6.769	59.899	1.192	34.760	5.009	0.590	1.928	0.300	0.002	0.543
18.387	0.141	43.565	24.420	0.433	7.004	31.764	1.005	1.147	0.219
144.695	2.663	17.967	0.091	9.003	3.217	14.382	1.008	2.336	4.562

- Construct a 95% CI and a 99% CI for  $\theta = \mathbb{E}[X]$ .

$n = 50$ ,  $\hat{\theta} = \bar{X} = 11.894$ ,  $S = 24.953$ . We use CI (4) and get  $t_{49, 0.975} = 2.010$ ,  $t_{49, 0.995} = 2.680$ . Then,

$$95\% \text{ CI: } 11.894 \pm 2.010 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 7.093 = [4.801, 18.987];$$

$$99\% \text{ CI: } 11.894 \pm 2.680 \times \frac{24.953}{\sqrt{50}} = 11.894 \pm 9.457 = [2.437, 21.351].$$

- Want to make half-length  $H \leq 2$  under 95% confidence level.

We use (6) and get  $z_{0.975} = 1.960$ ,  $S_0 = 24.953$ ,  $\epsilon = 2$ . Then,

$$n^* = \left\lceil \left( \frac{1.960 \times 24.953}{2} \right)^2 \right\rceil = \lceil 597.995 \rceil = 598.$$

Take  $598 - 50 = 548$  additional sample points.

- 1 Introduction
  - ▶ Types of Simulations
- 2 Point and Interval Estimations
  - ▶ Basics
  - ▶ Specified Precision
  - ▶ Example
- 3 Terminating Simulation**
  - ▶ Discrete Outputs**
  - ▶ Continuous Outputs**
- 4 Steady-State Simulation
  - ▶ Initialization Bias
  - ▶ Intelligent Initialization
  - ▶ Warm-up Period Deletion
  - ▶ Estimation with Multiple Replications
  - ▶ Estimation with Single Replication



# Terminating Simulation

- A terminating simulation runs over the time interval  $[0, T_E]$  and produce observations (outputs).

# Terminating Simulation

- A terminating simulation runs over the time interval  $[0, T_E]$  and produce observations (outputs).
- Discrete outputs:  $\{Y_1, Y_2, \dots, Y_n\}$ .
  - $n$  may be deterministic or random depending on how  $T_E$  is specified.
  - E.g., waiting time of all customers.
  - A common goal is to estimate  $\theta := \mathbb{E}[\frac{1}{n} \sum_{i=1}^n Y_i]$ , e.g., the *expectation* of the *average* waiting time.

# Terminating Simulation

- A terminating simulation runs over the time interval  $[0, T_E]$  and produce observations (outputs).
- Discrete outputs:  $\{Y_1, Y_2, \dots, Y_n\}$ .
  - $n$  may be deterministic or random depending on how  $T_E$  is specified.
  - E.g., waiting time of all customers.
  - A common goal is to estimate  $\theta := \mathbb{E}[\frac{1}{n} \sum_{i=1}^n Y_i]$ , e.g., the *expectation* of the *average* waiting time.
- Continuous outputs:  $\{Y(t) : 0 \leq t \leq T_E\}$ .
  - $T_E$  may be deterministic or random.
  - E.g., number of customer in the waiting line at time  $t$ ,  $0 \leq t \leq T_E$ .
  - A common goal is to estimate  $\theta := \mathbb{E}[\frac{1}{T_E} \int_0^{T_E} Y(t)dt]$ , e.g., the *expectation* of the *average* waiting line length.



# Terminating Simulation

- A terminating simulation runs over the time interval  $[0, T_E]$  and produce observations (outputs).
- Discrete outputs:  $\{Y_1, Y_2, \dots, Y_n\}$ .
  - $n$  may be deterministic or random depending on how  $T_E$  is specified.
  - E.g., waiting time of all customers.
  - A common goal is to estimate  $\theta := \mathbb{E}[\frac{1}{n} \sum_{i=1}^n Y_i]$ , e.g., the *expectation* of the *average* waiting time.
- Continuous outputs:  $\{Y(t) : 0 \leq t \leq T_E\}$ .
  - $T_E$  may be deterministic or random.
  - E.g., number of customer in the waiting line at time  $t$ ,  $0 \leq t \leq T_E$ .
  - A common goal is to estimate  $\theta := \mathbb{E}[\frac{1}{T_E} \int_0^{T_E} Y(t)dt]$ , e.g., the *expectation* of the *average* waiting line length.
- In general, independent replications (runs) are used, each with a different random number stream.



- Within-replication data vs. across-replication data:

Replication	Within-Rep Data (each row)	Across-Rep Data
1	$Y_{11}, Y_{12}, \dots, Y_{1n_1}$	$\bar{Y}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} Y_{1i}$
2	$Y_{21}, Y_{22}, \dots, Y_{2n_2}$	$\bar{Y}_2 = \frac{1}{n_2} \sum_{i=1}^{n_2} Y_{2i}$
$\vdots$	$\vdots$	$\vdots$
$R$	$Y_{R1}, Y_{R2}, \dots, Y_{Rn_R}$	$\bar{Y}_R = \frac{1}{n_R} \sum_{i=1}^{n_R} Y_{Ri}$

- Within-replication data vs. across-replication data:

Replication	Within-Rep Data (each row)	Across-Rep Data
1	$Y_{11}, Y_{12}, \dots, Y_{1n_1}$	$\bar{Y}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} Y_{1i}$
2	$Y_{21}, Y_{22}, \dots, Y_{2n_2}$	$\bar{Y}_2 = \frac{1}{n_2} \sum_{i=1}^{n_2} Y_{2i}$
$\vdots$	$\vdots$	$\vdots$
$R$	$Y_{R1}, Y_{R2}, \dots, Y_{Rn_R}$	$\bar{Y}_R = \frac{1}{n_R} \sum_{i=1}^{n_R} Y_{Ri}$

- Across-rep data are independent and identically distributed, when *same* initial conditions and *different* random number streams are used.

- **Within-replication** data vs. **across-replication** data:

Replication	Within-Rep Data (each row)	Across-Rep Data
1	$Y_{11}, Y_{12}, \dots, Y_{1n_1}$	$\bar{Y}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} Y_{1i}$
2	$Y_{21}, Y_{22}, \dots, Y_{2n_2}$	$\bar{Y}_2 = \frac{1}{n_2} \sum_{i=1}^{n_2} Y_{2i}$
$\vdots$	$\vdots$	$\vdots$
$R$	$Y_{R1}, Y_{R2}, \dots, Y_{Rn_R}$	$\bar{Y}_R = \frac{1}{n_R} \sum_{i=1}^{n_R} Y_{Ri}$

- Across-rep data are independent and identically distributed, when *same* initial conditions and *different* random number streams are used.
- Within-rep data are typically **neither independent nor identically distributed**:
  - waiting times of successive customers are heavily correlated;
  - waiting times during the peak hours are longer than off-peak hours, so they're not identically distributed.

- **Within-replication** data vs. **across-replication** data:

Replication	Within-Rep Data (each row)	Across-Rep Data
1	$Y_{11}, Y_{12}, \dots, Y_{1n_1}$	$\bar{Y}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} Y_{1i}$
2	$Y_{21}, Y_{22}, \dots, Y_{2n_2}$	$\bar{Y}_2 = \frac{1}{n_2} \sum_{i=1}^{n_2} Y_{2i}$
$\vdots$	$\vdots$	$\vdots$
$R$	$Y_{R1}, Y_{R2}, \dots, Y_{Rn_R}$	$\bar{Y}_R = \frac{1}{n_R} \sum_{i=1}^{n_R} Y_{Ri}$

- Across-rep data are independent and identically distributed, when *same* initial conditions and *different* random number streams are used.
- Within-rep data are typically **neither independent nor identically distributed**:
  - waiting times of successive customers are heavily correlated;
  - waiting times during the peak hours are longer than off-peak hours, so they're not identically distributed.
- Use **across-rep** data to do point/interval estimation!

- Example: What is the expectation of average waiting time for customers during  $[0, T_E]$ ?
  - Use  $\{\bar{Y}_1, \dots, \bar{Y}_R\}$  as an iid sample of size  $R$ .

- Example: What is the expectation of average waiting time for customers during  $[0, T_E]$ ?
  - Use  $\{\bar{Y}_1, \dots, \bar{Y}_R\}$  as an iid sample of size  $R$ .

- Point estimator:

$$\bar{Y} = \frac{1}{R} \sum_{r=1}^R \bar{Y}_r.$$

- Example: What is the expectation of average waiting time for customers during  $[0, T_E]$ ?
  - Use  $\{\bar{Y}_1, \dots, \bar{Y}_R\}$  as an iid sample of size  $R$ .

- Point estimator:

$$\bar{Y} = \frac{1}{R} \sum_{r=1}^R \bar{Y}_r.$$

- $1 - \alpha$  CI using (4):

$$\left[ \bar{Y} - t_{R-1, 1-\alpha/2} \frac{S}{\sqrt{R}}, \bar{Y} + t_{R-1, 1-\alpha/2} \frac{S}{\sqrt{R}} \right],$$

where  $S^2 = \frac{1}{R-1} \sum_{r=1}^R (\bar{Y}_r - \bar{Y})^2$ .

- Example: What is the expectation of average waiting time for customers during  $[0, T_E]$ ?
  - Use  $\{\bar{Y}_1, \dots, \bar{Y}_R\}$  as an iid sample of size  $R$ .

- Point estimator:

$$\bar{Y} = \frac{1}{R} \sum_{r=1}^R \bar{Y}_r.$$

- $1 - \alpha$  CI using (4):

$$\left[ \bar{Y} - t_{R-1, 1-\alpha/2} \frac{S}{\sqrt{R}}, \bar{Y} + t_{R-1, 1-\alpha/2} \frac{S}{\sqrt{R}} \right],$$

where  $S^2 = \frac{1}{R-1} \sum_{r=1}^R (\bar{Y}_r - \bar{Y})^2$ .

- Necessary number of replications for specified precision  $H \leq \epsilon$  under  $1 - \alpha$  confidence level, can be computed using (6).





- Within-replication data vs. across-replication data:

Replication	Within-Rep Data	Across-Rep Data
1	$\{Y_1(t) : 0 \leq t \leq T_{E_1}\}$	$\tilde{Y}_1 = \frac{1}{T_{E_1}} \int_0^{T_{E_1}} Y_1(t) dt$
2	$\{Y_2(t) : 0 \leq t \leq T_{E_2}\}$	$\tilde{Y}_2 = \frac{1}{T_{E_2}} \int_0^{T_{E_2}} Y_2(t) dt$
⋮	⋮	⋮
$R$	$\{Y_R(t) : 0 \leq t \leq T_{E_R}\}$	$\tilde{Y}_R = \frac{1}{T_{E_R}} \int_0^{T_{E_R}} Y_R(t) dt$

- Within-replication data vs. across-replication data:

Replication	Within-Rep Data	Across-Rep Data
1	$\{Y_1(t) : 0 \leq t \leq T_{E_1}\}$	$\tilde{Y}_1 = \frac{1}{T_{E_1}} \int_0^{T_{E_1}} Y_1(t) dt$
2	$\{Y_2(t) : 0 \leq t \leq T_{E_2}\}$	$\tilde{Y}_2 = \frac{1}{T_{E_2}} \int_0^{T_{E_2}} Y_2(t) dt$
$\vdots$	$\vdots$	$\vdots$
$R$	$\{Y_R(t) : 0 \leq t \leq T_{E_R}\}$	$\tilde{Y}_R = \frac{1}{T_{E_R}} \int_0^{T_{E_R}} Y_R(t) dt$

- Across-rep data are independent and identically distributed, when *same* initial conditions and *different* random number streams are used.

- Within-replication data vs. across-replication data:

Replication	Within-Rep Data	Across-Rep Data
1	$\{Y_1(t) : 0 \leq t \leq T_{E_1}\}$	$\tilde{Y}_1 = \frac{1}{T_{E_1}} \int_0^{T_{E_1}} Y_1(t) dt$
2	$\{Y_2(t) : 0 \leq t \leq T_{E_2}\}$	$\tilde{Y}_2 = \frac{1}{T_{E_2}} \int_0^{T_{E_2}} Y_2(t) dt$
⋮	⋮	⋮
$R$	$\{Y_R(t) : 0 \leq t \leq T_{E_R}\}$	$\tilde{Y}_R = \frac{1}{T_{E_R}} \int_0^{T_{E_R}} Y_R(t) dt$

- Across-rep data are independent and identically distributed, when *same* initial conditions and *different* random number streams are used.
- Example: What is the expectation of the average waiting line length during  $[0, T_E]$ ?
  - Use  $\{\tilde{Y}_1, \dots, \tilde{Y}_R\}$  as an iid sample of size  $R$ , and the rest is similar as before.

- 1 Introduction
  - ▶ Types of Simulations
- 2 Point and Interval Estimations
  - ▶ Basics
  - ▶ Specified Precision
  - ▶ Example
- 3 Terminating Simulation
  - ▶ Discrete Outputs
  - ▶ Continuous Outputs
- 4 Steady-State Simulation
  - ▶ Initialization Bias
  - ▶ Intelligent Initialization
  - ▶ Warm-up Period Deletion
  - ▶ Estimation with Multiple Replications
  - ▶ Estimation with Single Replication



# Steady-State Simulation

- Consider a single run of a simulation model whose purpose is to estimate a steady-state, or long-run, performance measure of the system.
  - Theoretically speaking, such steady-state performance measure has nothing to do with initial conditions.

# Steady-State Simulation

- Consider a single run of a simulation model whose purpose is to estimate a steady-state, or long-run, performance measure of the system.
  - Theoretically speaking, such steady-state performance measure has nothing to do with initial conditions.
- Discrete outputs:  $\{Y_1, Y_2, \dots\}$ .
  - A common goal is to estimate  $\phi := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n Y_i$ .

# Steady-State Simulation

- Consider a single run of a simulation model whose purpose is to estimate a steady-state, or long-run, performance measure of the system.
  - Theoretically speaking, such steady-state performance measure has nothing to do with initial conditions.
- Discrete outputs:  $\{Y_1, Y_2, \dots\}$ .
  - A common goal is to estimate  $\phi := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n Y_i$ .
- Continuous outputs:  $\{Y(t) : t \geq 0\}$ .
  - A common goal is to estimate  $\phi := \lim_{T_E \rightarrow \infty} \frac{1}{T_E} \int_0^{T_E} Y(t) dt$ .

# Steady-State Simulation

- Consider a single run of a simulation model whose purpose is to estimate a steady-state, or long-run, performance measure of the system.
  - Theoretically speaking, such steady-state performance measure has nothing to do with initial conditions.
- Discrete outputs:  $\{Y_1, Y_2, \dots\}$ .
  - A common goal is to estimate  $\phi := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n Y_i$ .
- Continuous outputs:  $\{Y(t) : t \geq 0\}$ .
  - A common goal is to estimate  $\phi := \lim_{T_E \rightarrow \infty} \frac{1}{T_E} \int_0^{T_E} Y(t) dt$ .
- However, we cannot simulate a system “to infinity” but must stop somewhere.
  - The simulation run length ( $n$  or  $T_E$ ) is a design choice instead of inherently determined by the nature of the problem.



- The run length in steady-state simulation needs to be *carefully chosen*, with several considerations:
  - bias that is due to artificial or arbitrary initial conditions;
    - can be severe if run length is too short
    - generally decreases as run length increases

- The run length in steady-state simulation needs to be *carefully chosen*, with several considerations:
  - bias that is due to artificial or arbitrary initial conditions;
    - can be severe if run length is too short
    - generally decreases as run length increases
  - the desired precision of the point estimator;
    - measured by the **standard error** (i.e., estimator's standard deviation) or confidence interval half-width

- The run length in steady-state simulation needs to be *carefully chosen*, with several considerations:
  - bias that is due to artificial or arbitrary initial conditions;
    - can be severe if run length is too short
    - generally decreases as run length increases
  - the desired precision of the point estimator;
    - measured by the **standard error** (i.e., estimator's standard deviation) or confidence interval half-width
  - budget constraints on the time available to execute the simulation.

- The effect of the initial condition is persistent and typically does not vanish after a *finite* time.
  - Unless the initial condition is specified as the “stationary distribution” of the system, which is unknown in general.

- The effect of the initial condition is persistent and typically does not vanish after a *finite* time.
  - Unless the initial condition is specified as the “stationary distribution” of the system, which is unknown in general.
- **Fact 1:** Estimators based on a finite-time simulation (finite  $n$  or  $T_E$ ) are biased:

$$\mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n Y_i \right] \neq \phi, \quad \mathbb{E} \left[ \frac{1}{T_E} \int_0^{T_E} Y(t) dt \right] \neq \phi.$$

- The effect of the initial condition is persistent and typically does not vanish after a *finite* time.
  - Unless the initial condition is specified as the “stationary distribution” of the system, which is unknown in general.
- **Fact 1:** Estimators based on a finite-time simulation (finite  $n$  or  $T_E$ ) are biased:

$$\mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n Y_i \right] \neq \phi, \quad \mathbb{E} \left[ \frac{1}{T_E} \int_0^{T_E} Y(t) dt \right] \neq \phi.$$

- **Fact 2:** The bias cannot be replicated away:

$$\lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=1}^R \bar{Y}_r \neq \phi, \quad \lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=1}^R \tilde{Y}_r \neq \phi.$$

- The effect of the initial condition is persistent and typically does not vanish after a *finite* time.
  - Unless the initial condition is specified as the “stationary distribution” of the system, which is unknown in general.
- **Fact 1:** Estimators based on a finite-time simulation (finite  $n$  or  $T_E$ ) are biased:

$$\mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n Y_i \right] \neq \phi, \quad \mathbb{E} \left[ \frac{1}{T_E} \int_0^{T_E} Y(t) dt \right] \neq \phi.$$

- **Fact 2:** The bias cannot be replicated away:

$$\lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=1}^R \bar{Y}_r \neq \phi, \quad \lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=1}^R \tilde{Y}_r \neq \phi.$$

- With more replications, we get a more “precise” estimate of an incorrect value.
  - The confidence interval is narrower but it is centered at an incorrect position.

- Example of  $M/M/1$  queue: <https://xiaoweiz.shinyapps.io/MM1queue>
  - If  $\lambda < \mu$ , the system is stable and the **steady-state expectation** (or **long-run average**) of waiting time is

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n Y_i = \frac{\lambda/\mu}{\mu - \lambda}.$$



- Example of  $M/M/1$  queue: <https://xiaoweiz.shinyapps.io/MM1queue>
  - If  $\lambda < \mu$ , the system is stable and the **steady-state expectation** (or **long-run average**) of waiting time is

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n Y_i = \frac{\lambda/\mu}{\mu - \lambda}.$$

- Choosing different initial conditions (in this example, number of customers in station, also known as initial state) gives different looks of sample paths (over finite time period).

- Example of  $M/M/1$  queue: <https://xiaoweiz.shinyapps.io/MM1queue>
  - If  $\lambda < \mu$ , the system is stable and the **steady-state expectation** (or **long-run average**) of waiting time is

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n Y_i = \frac{\lambda/\mu}{\mu - \lambda}.$$

- Choosing different initial conditions (in this example, number of customers in station, also known as initial state) gives different looks of sample paths (over finite time period).
- Methods to reduce initialization bias:
  - intelligent initialization;
  - warm-up period deletion;
  - low-bias estimator (advanced topic).

- Initialize the simulation in a state that is more representative of long-run conditions.

- Initialize the simulation in a state that is more representative of long-run conditions.
- If the system exists, collect data on it and use these data to specify more nearly typical initial conditions:
  - fit a probability distribution to describe the initial state;
  - or, simply use the sample mean as a representative.

- Initialize the simulation in a state that is more representative of long-run conditions.
- If the system exists, collect data on it and use these data to specify more nearly typical initial conditions:
  - fit a probability distribution to describe the initial state;
  - or, simply use the sample mean as a representative.
- If the system can be simplified enough to make it analytically solvable, e.g. queueing models, use the theoretical solution to initialize the simulation.
  - Solve the simplified model to find the stationary distribution or most likely conditions (e.g., the expected number of customers in a station).
  - This is another important value of those analytically solvable queueing models.

- The impact of the initial condition gradually vanishes as the run length increases.

- The impact of the initial condition gradually vanishes as the run length increases.
- So we divide a simulation run into two periods:
  - warm-up period: from time 0 to time  $T_0$ ;
  - data-collection period: from time  $T_0$  to  $T_E$ .

- The impact of the initial condition gradually vanishes as the run length increases.
- So we divide a simulation run into two periods:
  - warm-up period: from time 0 to time  $T_0$ ;
  - data-collection period: from time  $T_0$  to  $T_E$ .




Figure: Warm-up Period Deletion (from [Banks et al. \(2010\)](#))



- The impact of the initial condition gradually vanishes as the run length increases.
- So we divide a simulation run into two periods:
  - warm-up period: from time 0 to time  $T_0$ ;
  - data-collection period: from time  $T_0$  to  $T_E$ .




Figure: Warm-up Period Deletion (from [Banks et al. \(2010\)](#))

- $T_0$  should be sufficiently large so that at time  $T_0$  the impact of the initial condition is very weak and the system behaves approximately as in the steady state.

- To determine  $T_0$ 
  - There are no widely accepted and proven techniques.
  - Plots are often used.
- The raw output data plot from a single simulation run is usually too fluctuating to detect the trend. – **not helpful**
- Instead of directly plotting raw output data, we usually use some smoother plots to see when the curve “stabilizes”:
  - cumulative average (累积均值); – **OK**
  - ensemble average (总体均值). – **recommended**




Figure: Raw Output of Waiting Time of Each Customer in  $M/M/1$  Queue with  $\lambda = 0.962$  and  $\mu = 1$  (from [ZHANG Xiaowei](#))

- Cumulative average (累积均值): For one replication, say, replication 1, plot the average from time 0 up to now.
  - Discrete outputs: Plot  $\bar{Y}_1(n) = \frac{1}{n} \sum_{i=1}^n Y_{1i}$  with respect to  $n$ ;
  - Continuous outputs: Plot  $\tilde{Y}_1(T) = \frac{1}{T} \int_0^T Y_1(t) dt$  with respect to  $T$ .

- Cumulative average (累积均值): For one replication, say, replication 1, plot the average from time 0 up to now.
  - Discrete outputs: Plot  $\bar{Y}_1(n) = \frac{1}{n} \sum_{i=1}^n Y_{1i}$  with respect to  $n$ ;
  - Continuous outputs: Plot  $\tilde{Y}_1(T) = \frac{1}{T} \int_0^T Y_1(t) dt$  with respect to  $T$ .
- It can be plotted for each replication, so we usually detect different warm-up period durations from different replications.

- Cumulative average (累积均值): For one replication, say, replication 1, plot the average from time 0 up to now.
  - Discrete outputs: Plot  $\bar{Y}_1(n) = \frac{1}{n} \sum_{i=1}^n Y_{1i}$  with respect to  $n$ ;
  - Continuous outputs: Plot  $\tilde{Y}_1(T) = \frac{1}{T} \int_0^T Y_1(t) dt$  with respect to  $T$ .
- It can be plotted for each replication, so we usually detect different warm-up period durations from different replications.
- The cumulative plot is usually conservative, i.e., the warm-up period it detects is longer than necessary.
  - It retains all of the data including the warm-up period, so the bias needs more time to diminish.




Figure: Cumulative Average Waiting Time of Customers in  $M/M/1$  Queue with  $\lambda = 0.962$  and  $\mu = 1$  (from [ZHANG Xiaowei](#))

- Ensemble average (总体均值): For multiple replications  $1, \dots, R$ , compute the average across replications and make the plot.
  - Discrete outputs: Plot  $\bar{Y}(n) = \frac{1}{R} \sum_{r=1}^R Y_{nr}$  with respect to  $n$ ;
  - Continuous outputs: *Divide the raw data of replication  $r$  into small batches*, e.g.,  $\{Y_r(t) : (j-1)m \leq t < jm\}$ ,  $j = 1, 2, \dots$ ;  
plot  $\tilde{Y}(j) = \frac{1}{R} \sum_{r=1}^R \left[ \frac{1}{m} \int_{(j-1)m}^{jm} Y_r(t) dt \right]$  with respect to  $j$ .



- Ensemble average (总体均值): For multiple replications  $1, \dots, R$ , compute the average across replications and make the plot.
  - Discrete outputs: Plot  $\bar{Y}(n) = \frac{1}{R} \sum_{r=1}^R Y_{nr}$  with respect to  $n$ ;
  - Continuous outputs: *Divide the raw data of replication  $r$  into small batches*, e.g.,  $\{Y_r(t) : (j-1)m \leq t < jm\}$ ,  $j = 1, 2, \dots$ ;  
plot  $\tilde{Y}(j) = \frac{1}{R} \sum_{r=1}^R \left[ \frac{1}{m} \int_{(j-1)m}^{jm} Y_r(t) dt \right]$  with respect to  $j$ .
- We detect one warm-up period duration for multiple replications.

- Ensemble average (总体均值): For multiple replications  $1, \dots, R$ , compute the average across replications and make the plot.
  - Discrete outputs: Plot  $\bar{Y}(n) = \frac{1}{R} \sum_{r=1}^R Y_{nr}$  with respect to  $n$ ;
  - Continuous outputs: *Divide the raw data of replication  $r$  into small batches*, e.g.,  $\{Y_r(t) : (j-1)m \leq t < jm\}$ ,  $j = 1, 2, \dots$ ;  
plot  $\tilde{Y}(j) = \frac{1}{R} \sum_{r=1}^R \left[ \frac{1}{m} \int_{(j-1)m}^{jm} Y_r(t) dt \right]$  with respect to  $j$ .
- We detect one warm-up period duration for multiple replications.
- Some variations are smoothed out by averaging across multiple replications.
  - This leads to more accurate detection of warm-up period.




Figure: Ensemble Average Waiting Time of  $n$ -th Customer in  $M/M/1$  Queue with  $\lambda = 0.962$  and  $\mu = 1$  (from [ZHANG Xiaowei](#))

- When first starting to detect the warm-up period, a run length and number of replications will have to be guessed.
  - Increase the number of replications if the ensemble averages are not smooth enough.
  - Increase the run length if the ensemble averages do not stabilize.

- When first starting to detect the warm-up period, a run length and number of replications will have to be guessed.
  - Increase the number of replications if the ensemble averages are not smooth enough.
  - Increase the run length if the ensemble averages do not stabilize.
- Since each ensemble average is the sample mean of iid observations across  $R$  replications, a confidence interval can be placed around each point.
  - Use them to judge whether or not the plot is precise enough to decide that the bias has vanished.
  - This is the preferred method to determine a deletion point.




Figure: Ensemble Average Waiting Time and 95% CI of  $n$ -th Customer in  $M/M/1$  Queue with  $\lambda = 0.962$  and  $\mu = 1$  (from [ZHANG Xiaowei](#))

- Cumulative averages become less variable as more data are averaged.
  - So the right side of the curve is always smoother than the left side.

- Cumulative averages become less variable as more data are averaged.
  - So the right side of the curve is always smoother than the left side.
- Cumulative averages tend to converge more slowly to long-run performance than ensemble averages do.
  - Because cumulative averages contain all observations including the most biased ones from the very beginning.
  - Cumulative averages should be used *only if* ensemble averages can not be computed, such as when only a single replication is possible.



- Cumulative averages become less variable as more data are averaged.
  - So the right side of the curve is always smoother than the left side.
- Cumulative averages tend to converge more slowly to long-run performance than ensemble averages do.
  - Because cumulative averages contain all observations including the most biased ones from the very beginning.
  - Cumulative averages should be used *only if* ensemble averages can not be computed, such as when only a single replication is possible.
- Different performance measures could approach steady state with different speed.

- **Idea:** Make multiple replications (long enough), remove warm-up period for each one, and then work as if we were in a terminating simulation.

- **Idea:** Make multiple replications (long enough), remove warm-up period for each one, and then work as if we were in a terminating simulation.
- **Caution:** Make sure that initialization bias in the point estimator has been reduced to a negligible level.
  - Otherwise the estimation can be misleading.

- **Idea:** Make multiple replications (long enough), remove warm-up period for each one, and then work as if we were in a terminating simulation.
- **Caution:** Make sure that initialization bias in the point estimator has been reduced to a negligible level.
  - Otherwise the estimation can be misleading.
- **Note:** Initialization bias is not affected by the number of replications.
  - It is affected by deleting more data (i.e. increasing  $T_0$ ) or extending the run length (i.e. increasing  $T_E$ ).
  - Increasing the number of replications could produce narrower interval around the “wrong point”.

- Discrete outputs:
  - Suppose we decide to delete first  $d$  observations of the total  $n$  observations in a replication.<sup>†</sup>
  - The across-replication data from  $R$  replications are

$$\bar{Y}_1 = \frac{1}{n-d} \sum_{i=d+1}^n Y_{1i}, \dots, \bar{Y}_R = \frac{1}{n-d} \sum_{i=d+1}^n Y_{Ri}.$$

---

<sup>†</sup>  $d$  and  $n$  may vary between different replications, in which case they are replaced by  $d_r$  and  $n_r$ , respectively.

- Discrete outputs:
  - Suppose we decide to delete first  $d$  observations of the total  $n$  observations in a replication.<sup>†</sup>
  - The across-replication data from  $R$  replications are

$$\bar{Y}_1 = \frac{1}{n-d} \sum_{i=d+1}^n Y_{1i}, \dots, \bar{Y}_R = \frac{1}{n-d} \sum_{i=d+1}^n Y_{Ri}.$$

- Continuous outputs:
  - Suppose we decide to delete data in  $[0, T_0]$  period and only use those in  $[T_0, T_E]$  in a replication.
  - The across-replication data from  $R$  replications are

$$\tilde{Y}_1 = \frac{1}{T_E - T_0} \int_{T_0}^{T_E} Y_1(t) dt, \dots, \tilde{Y}_R = \frac{1}{T_E - T_0} \int_{T_0}^{T_E} Y_R(t) dt.$$

---

<sup>†</sup>  $d$  and  $n$  may vary between different replications, in which case they are replaced by  $d_r$  and  $n_r$ , respectively.

- Similar as in terminating simulation, the across-replication data are iid.
  - So we can use them to compute the point estimator, CI, and necessary number of replications for specified precision, in the same way as before.

- Similar as in terminating simulation, the across-replication data are iid.
  - So we can use them to compute the point estimator, CI, and necessary number of replications for specified precision, in the same way as before.
- Unlike terminating simulation, the above mentioned estimators are biased for finite  $n$  or  $T_E$ .
  - The bias is negligible if  $d$  and  $n$ , or  $T_0$  and  $T_E$ , are sufficiently large.



- Similar as in terminating simulation, the across-replication data are iid.
  - So we can use them to compute the point estimator, CI, and necessary number of replications for specified precision, in the same way as before.
- Unlike terminating simulation, the above mentioned estimators are biased for finite  $n$  or  $T_E$ .
  - The bias is negligible if  $d$  and  $n$ , or  $T_0$  and  $T_E$ , are sufficiently large.
- A rough rule for relationship between  $d$  and  $n$ , or  $T_0$  and  $T_E$ :

$$(n - d) \geq 10d, \quad (T_E - T_0) \geq 10T_0.$$

- Suppose analysis indicates that  $R - R_0$  additional replications are needed after the initial number  $R_0$ , in order to meet the desired precision.
- An alternative to increasing replications is to increase run length  $T_E$  within each replication.

- Suppose analysis indicates that  $R - R_0$  additional replications are needed after the initial number  $R_0$ , in order to meet the desired precision.
- An alternative to increasing replications is to increase run length  $T_E$  within each replication.
  - Increase run length  $T_E$  in the same proportion  $(R/R_0)$  to a new run length  $(R/R_0)T_E$ .
  - More data will be deleted, from time 0 to time  $(R/R_0)T_0$ .
  - More data will be used to compute the estimate, from time  $(R/R_0)T_0$  to time  $(R/R_0)T_E$ .
  - The total amount of simulation effort is the same as if we had simply increased the number of replications.




Figure: Increasing Run Length to Achieve Specified Precision (from [Banks et al. \(2010\)](#))




Figure: Increasing Run Length to Achieve Specified Precision (from [Banks et al. \(2010\)](#))

- Advantage: Any residual bias in the point estimator would be further reduced.




Figure: Increasing Run Length to Achieve Specified Precision (from [Banks et al. \(2010\)](#))

- Advantage: Any residual bias in the point estimator would be further reduced.
- Disadvantage: It is necessary to have saved the state of the model at time  $T_E$  and to be able to continue the running.
  - Otherwise, the simulations would have to be re-run from time 0, which could be time consuming for a complex model.

- A disadvantage of the replication method is that the warm-up period must be deleted on each replication.
  - This can become very costly in terms of computation time especially when the model warms up very slowly.
    - E.g.,  $M/M/1$  queue with utilization close to 1.

- A disadvantage of the replication method is that the warm-up period must be deleted on each replication.
  - This can become very costly in terms of computation time especially when the model warms up very slowly.
    - E.g.,  $M/M/1$  queue with utilization close to 1.
- This suggests that we could use one single, (very) long replication for estimation, so that only one warm-up period is deleted.
- Besides, it is also possible that we are in a situation where only the data from one long replication are available.



- Point estimator: Sample mean after the warm-up period deletion

$$\bar{Y} = \frac{1}{n-d} \sum_{i=d+1}^n Y_i, \quad \tilde{Y} = \frac{1}{T_E - T_0} \int_{T_0}^{T_E} Y(t) dt.$$

- Point estimator: Sample mean after the warm-up period deletion

$$\bar{Y} = \frac{1}{n-d} \sum_{i=d+1}^n Y_i, \quad \tilde{Y} = \frac{1}{T_E - T_0} \int_{T_0}^{T_E} Y(t) dt.$$

- The disadvantage of the single-replication design arises when we try to estimate the variance of the above estimators, because of
  - the strong but unknown dependence among  $Y_1, Y_2, \dots, Y_n$ ;
  - the non-identical distribution of  $Y_1, Y_2, \dots, Y_n$ ;
  - and the integral form of  $\tilde{Y}$ .

- **Caution:** It is tempting to compute

$$S^2 = \frac{1}{n-d-1} \sum_{i=d+1}^n (Y_i - \bar{Y})^2,$$

and use  $S^2/(n-d)$  to estimate  $\text{Var}(\bar{Y})$ . However, such estimation will be **terrible**, since  $Y_1, Y_2, \dots, Y_n$  are **neither independent nor identically distributed**.

- **Caution:** It is tempting to compute

$$S^2 = \frac{1}{n-d-1} \sum_{i=d+1}^n (Y_i - \bar{Y})^2,$$

and use  $S^2/(n-d)$  to estimate  $\text{Var}(\bar{Y})$ . However, such estimation will be **terrible**, since  $Y_1, Y_2, \dots, Y_n$  are **neither independent nor identically distributed**.

- The CI based on  $S^2/(n-d)$  would also be misleading.

- **Caution:** It is tempting to compute

$$S^2 = \frac{1}{n-d-1} \sum_{i=d+1}^n (Y_i - \bar{Y})^2,$$

and use  $S^2/(n-d)$  to estimate  $\text{Var}(\bar{Y})$ . However, such estimation will be **terrible**, since  $Y_1, Y_2, \dots, Y_n$  are **neither independent nor identically distributed**.

- The CI based on  $S^2/(n-d)$  would also be misleading.
- Example: Suppose  $Y_{d+1}, \dots, Y_n$  are identically distributed but positive correlated (which is common for waiting time), then

$$\mathbb{E}[S^2/(n-d)] < \text{Var}(\bar{Y}).$$

- **Caution:** It is tempting to compute

$$S^2 = \frac{1}{n-d-1} \sum_{i=d+1}^n (Y_i - \bar{Y})^2,$$

and use  $S^2/(n-d)$  to estimate  $\text{Var}(\bar{Y})$ . However, such estimation will be **terrible**, since  $Y_1, Y_2, \dots, Y_n$  are **neither independent nor identically distributed**.

- The CI based on  $S^2/(n-d)$  would also be misleading.
- Example: Suppose  $Y_{d+1}, \dots, Y_n$  are identically distributed but positive correlated (which is common for waiting time), then

$$\mathbb{E}[S^2/(n-d)] < \text{Var}(\bar{Y}).$$

- The constructed CI using  $S^2/(n-d)$  will be narrower than the actual valid one.

- Batch Means (批均值) Method:
  - Divide the output data from one replication (after deleting warm-up period) into  $k$  **large batches**, and compute the batch means.

- Batch Means (批均值) Method:
  - Divide the output data from one replication (after deleting warm-up period) into  $k$  **large batches**, and compute the batch means.
  - Discrete outputs:  $\bar{Y}_j = \frac{1}{m} \sum_{i=(j-1)m+1}^{jm} Y_{i+d}$ ,  $j = 1, \dots, k$ .

$$\underbrace{Y_1, \dots, Y_d}_{\text{deleted}}, \underbrace{Y_{d+1}, \dots, Y_{d+m}}_{\text{Batch 1: } \bar{Y}_1}, \underbrace{Y_{d+m+1}, \dots, Y_{d+2m}}_{\text{Batch 2: } \bar{Y}_2}, \dots, \underbrace{Y_{d+(k-1)m+1}, \dots, Y_{d+km}}_{\text{Batch } k: \bar{Y}_k}$$



- Batch Means (批均值) Method:
  - Divide the output data from one replication (after deleting warm-up period) into  $k$  **large batches**, and compute the batch means.

- Discrete outputs:  $\bar{Y}_j = \frac{1}{m} \sum_{i=(j-1)m+1}^{jm} Y_{i+d}$ ,  $j = 1, \dots, k$ .

$$\underbrace{Y_1, \dots, Y_d}_{\text{deleted}}, \underbrace{Y_{d+1}, \dots, Y_{d+m}}_{\text{Batch 1: } \bar{Y}_1}, \underbrace{Y_{d+m+1}, \dots, Y_{d+2m}}_{\text{Batch 2: } \bar{Y}_2}, \dots, \underbrace{Y_{d+(k-1)m+1}, \dots, Y_{d+km}}_{\text{Batch } k: \bar{Y}_k}$$

- Continuous outputs:  $\tilde{Y}_j = \frac{1}{m} \int_{T_0+(j-1)m}^{T_0+jm} Y(t) dt$ ,  $j = 1, \dots, k$ .

- Batch Means (批均值) Method:

- Divide the output data from one replication (after deleting warm-up period) into  $k$  **large batches**, and compute the batch means.

- Discrete outputs:  $\bar{Y}_j = \frac{1}{m} \sum_{i=(j-1)m+1}^{jm} Y_{i+d}$ ,  $j = 1, \dots, k$ .

$$\underbrace{Y_1, \dots, Y_d}_{\text{deleted}}, \underbrace{Y_{d+1}, \dots, Y_{d+m}}_{\text{Batch 1: } \bar{Y}_1}, \underbrace{Y_{d+m+1}, \dots, Y_{d+2m}}_{\text{Batch 2: } \bar{Y}_2}, \dots, \underbrace{Y_{d+(k-1)m+1}, \dots, Y_{d+km}}_{\text{Batch } k: \bar{Y}_k}$$

- Continuous outputs:  $\tilde{Y}_j = \frac{1}{m} \int_{T_0+(j-1)m}^{T_0+jm} Y(t) dt$ ,  $j = 1, \dots, k$ .
- Treat the means of these batches **as if** they were independent.

- Batch Means (批均值) Method:
  - Divide the output data from one replication (after deleting warm-up period) into  $k$  **large batches**, and compute the batch means.

- Discrete outputs:  $\bar{Y}_j = \frac{1}{m} \sum_{i=(j-1)m+1}^{jm} Y_{i+d}$ ,  $j = 1, \dots, k$ .

$$\underbrace{Y_1, \dots, Y_d}_{\text{deleted}}, \underbrace{Y_{d+1}, \dots, Y_{d+m}}_{\text{Batch 1: } \bar{Y}_1}, \underbrace{Y_{d+m+1}, \dots, Y_{d+2m}}_{\text{Batch 2: } \bar{Y}_2}, \dots, \underbrace{Y_{d+(k-1)m+1}, \dots, Y_{d+km}}_{\text{Batch } k: \bar{Y}_k}$$

- Continuous outputs:  $\tilde{Y}_j = \frac{1}{m} \int_{T_0+(j-1)m}^{T_0+jm} Y(t)dt$ ,  $j = 1, \dots, k$ .
  - Treat the means of these batches **as if** they were independent.
- Why it works?
    - The correlation between two observations decreases as they are farther apart.
    - If the batch size is sufficiently large,
      - most of the observations in a batch will be approximately independent of those in other batches;
      - only those near the end of the batches are significantly correlated.

- Strictly speaking, the batch means are not independent.
- However, if the batch size is sufficiently large, successive batch means will be approximately independent.

- Strictly speaking, the batch means are not independent.
- However, if the batch size is sufficiently large, successive batch means will be approximately independent.
- Unfortunately, there is no widely accepted and relatively simple method for choosing an acceptable batch size  $m$  (or equivalently, choosing a number of batches  $k$ ).

- Strictly speaking, the batch means are not independent.
- However, if the batch size is sufficiently large, successive batch means will be approximately independent.
- Unfortunately, there is no widely accepted and relatively simple method for choosing an acceptable batch size  $m$  (or equivalently, choosing a number of batches  $k$ ).
- Some general guidelines:
  - In most applications, it is suggested to let  $10 \leq k \leq 30$ , according to [Schmeiser \(1982\)](#).
  - If the run length is to be increased to attain a specified precision, it is suggested to allow both  $m$  and  $k$  to grow.